

Vaccine	Vaccine description	ACIP Abbreviation	Trade Name	Age Limitation MIN	Age Limitation MAX
Coronavirus	COVID-19 mRNA monovalent	COVID-19	Moderna COVID-19	≥ 6 months old	none
	COVID-19 mRNA bivalent	COVID-19	Moderna COVID-19 Bivalent	≥ 6 months old	none
	COVID-19 mRNA monovalent	COVID-19	Pfizer-BioNTech COVID-19	≥ 6 months old	none
	COVID-19 mRNA bivalent	COVID-19	Pfizer-BioNTech COVID-19 Bivalent	≥ 6 months old	none
	SARS-CoV-2 recombinant spike protein adjuvant vaccine	COVID-19	Novavax COVID-19	≥ 12 years old	none
	recombinant, replication-incompetent adenovirus serotype 26 (Ad26) vector vaccine	COVID-19	Janssen COVID-19 vaccine	≥ 18 years	none
Diphtheria, tetanus and pertussis	Diphtheria and tetanus toxoids adsorbed	DT	Diphtheria and tetanus toxoids adsorbed	≥ 1 month* (6 weeks)	<7 years
	Diphtheria and tetanus toxoids and acellular pertussis vaccine adsorbed	DTaP	Daptacel, Infanrix	≥ 1 month* (6 weeks)	<7 years
	Tetanus and diphtheria toxoids adsorbed (A)	Td	Tenivac	≥ 7 years	none
	Tetanus toxoid, reduced diphtheria toxoid and acellular pertussis vaccine, adsorbed	Tdap	Adacel, Boostrix	≥ 7 years	none
	Diphtheria and tetanus toxoids and acellular pertussis adsorbed, hepatitis B and inactivated poliovirus vaccine	DTaP-HepB- IPV	Pediarix	≥ 1 month* (6 weeks)	<7 years

^{*}This reference chart is intended to provide vaccine abbreviations and age limitations used in ACIP Recommendations and Policy Notes for routine immunizations (non-routine immunization criteria are not included in this document); published in the *MMWR* and in the U.S. immunization schedules for children, adolescents, and adults. This information is subject to change.



Vaccine	Vaccine description	ACIP Abbreviation	Trade Name	Age Limitation MIN	Age Limitation MAX
	Diphtheria and tetanus toxoids and acellular pertussis adsorbed and inactivated poliovirus vaccine	DTaP-IPV	Kinrix Quadracel	≥ 4 years	<7 years
	Diphtheria and tetanus toxoids and acellular pertussis adsorbed, inactivated poliovirus and Haemophilus influenzae type b conjugate vaccine	DTaP-IPV/Hib	Pentacel, Vaxelis	≥ 1 month* (6 weeks)	<5 years
Haemophilus influenzae type b	Haemophilus influenzae type b conjugate vaccine	Hib	PedvaxHIB; ActHIB Hiberix	≥ 1 month* (6 weeks)	none
	H. influenzae type b, diphtheria, tetanus, pertussis, and polio vaccine	DTaP-IPV/Hib	Pentacel, Vaxelis	≥ 1 month* (6 weeks)	<5 years
Hepatitis	Hepatitis A vaccine	НерА	Havrix Vaqta	≥ 6 months	none
	Hepatitis B vaccine	НерВ	Engerix-B Recombivax HB	none	none
	Hepatitis B-TLR	HepB-CpG	Heplisav-B	≥ 18 years	none
	Hepatitis A inactivated and hepatitis B vaccine	НерА-НерВ	Twinrix	≥ 18 year	none
	Diphtheria and tetanus toxoids and acellular pertussis adsorbed,	DTaP-HepB- IPV	Pediarix	≥ 1 month* (6 weeks)	<7 years

^{*}This reference chart is intended to provide vaccine abbreviations and age limitations used in ACIP Recommendations and Policy Notes for routine immunizations (non-routine immunization criteria are not included in this document); published in the *MMWR* and in the U.S. immunization schedules for children, adolescents, and adults. This information is subject to change.



Vaccine	Vaccine description	ACIP Abbreviation	Trade Name	Age Limitation MIN	Age Limitation MAX
	hepatitis B and inactivated poliovirus vaccine				
Herpes zoster (shingles)	Zoster Vaccine Live	ZVL	Zostavax	≥ 60 years	none
	Zoster Vaccine Recombinant, Adjuvanted	RZV	Shingrix	≥ 19 years	none
Human papillomavirus	Human papillomavirus vaccine (9-valent) – types 6,11,16,18,31, 33,45,52,and 58	9vHPV	Gardasil-9	≥9 years	<46 years
Influenza	Live Attenuated Influenza Vaccine, quadrivalent	LAIV4	Flumist	≥ 2 years	<50 years
	Trivalent inactivated influenza vaccine	IIV3	Fluad	≥ 65 years	None
	Quadrivalent inactivated influenza vaccine	IIV4	Afluria Quad	≥ 6 months	None
			Fluad Quad	≥ 65 years	None
			Flublok Quad	≥ 18 years	None
			Fluarix Quad	≥ 6 months	None
			FluLaval Quad	≥ 6 months	None
			Flucelvax Quad	≥ 6 months	None
			Fluzone High Dose Quad	≥ 65 years	None

^{*}This reference chart is intended to provide vaccine abbreviations and age limitations used in ACIP Recommendations and Policy Notes for routine immunizations (non-routine immunization criteria are not included in this document); published in the *MMWR* and in the U.S. immunization schedules for children, adolescents, and adults. This information is subject to change.



Vaccine	Vaccine description	ACIP Abbreviation	Trade Name	Age Limitation MIN	Age Limitation MAX
			Fluzone Quad – MDV	≥ 6 months	None
			Fluzone Quad (Syringe/SDV)	≥ 6 months	None
	Recombinant Influenza Vaccine, Trivalent	RIV3	FluBlok	≥ 18 years	None
Measles, mumps, rubella and varicella	Measles, mumps, and rubella vaccine	MMR	M-M-R II Priorix	≥ 6 months	None
	Measles, mumps, rubella, and varicella vaccine	MMRV	ProQuad	≥1 year	None
Meningococcal	Quadrivalent	MenACWY- CRM	Menveo	≥2 months	None
	Quadrivalent	MenACWY- D	Menactra	≥9 months	None
	Quadrivalent tetanus toxoid conjugate vaccine	MenACWY-TT	MenQuadfi	≥2 years	None
	Serogroup B meningococcal vaccine	MenB-FHbp MenB-4C	Trumenba Bexsero	≥10 years	none
Pneumococcal	Pneumococcal conjugate vaccine (13-valent)	PCV13	Prevnar 13	≥ 1 month* (6 weeks)	none
	Pneumococcal polysaccharide vaccine (23-valent)	PPSV23	Pneumovax	≥ 2 years	none
	Pneumococcal conjugate vaccine (15-valent)	PCV15	Vaxneuvance	≥ 1 month* (6 weeks)	none

^{*}This reference chart is intended to provide vaccine abbreviations and age limitations used in ACIP Recommendations and Policy Notes for routine immunizations (non-routine immunization criteria are not included in this document); published in the *MMWR* and in the U.S. immunization schedules for children, adolescents, and adults. This information is subject to change.



Vaccine	Vaccine description	ACIP Abbreviation	Trade Name	Age Limitation MIN	Age Limitation MAX
	Pneumococcal conjugate vaccine (20-valent)	PCV20	Prevnar 20	≥ 19 years	none
Poliovirus	Inactivated poliovirus vaccine	IPV	lpol	≥ 1 month* (6 weeks)	none
	Diphtheria and tetanus toxoids and acellular pertussis adsorbed and inactivated poliovirus vaccine	DTaP-IPV	Kinrix Quadracel	≥ 4 years	<7 years
	Diphtheria and tetanus toxoids and acellular pertussis adsorbed, inactivated poliovirus and Haemophilus influenzae type b conjugate vaccine	DTaP-IPV/Hib	Pentacel, Vaxelis	≥ 1 month* (6 weeks)	<5 years
Rotavirus	Rotavirus vaccine (monovalent)	RV1	Rotarix	≥ 1 month* (6 weeks)	<9 months
	Rotavirus vaccine (pentavalent)	RV5	RotaTeq	≥ 1 month* (6 weeks)	<9 months
Varicella	Varicella vaccine	VAR	Varivax	≥ 1 years	none

^{*}This reference chart is intended to provide vaccine abbreviations and age limitations used in ACIP Recommendations and Policy Notes for routine immunizations (non-routine immunization criteria are not included in this document); published in the *MMWR* and in the U.S. immunization schedules for children, adolescents, and adults. This information is subject to change.